

DUISTERMAAT-HECKMAN FORMULA FOR A TORUS ACTION ON A GENERALIZED CALABI-YAU MANIFOLD AND LOCALIZATION FORMULA

YASUFUMI NITTA

ABSTRACT. This note is an addendum to our earlier work [6]. In [6], we studied a Hamiltonian action for a generalized Calabi-Yau manifold and showed that the Duistermaat-Heckman theorem holds. The purpose of this note is to show that the density function of the Duistermaat-Heckman measure is a piecewise polynomial. We also prove that the localization formula holds.

1. INTRODUCTION

In [6], we studied a Hamiltonian action for a generalized Calabi-Yau manifold. Let M be a $2n$ -dimensional connected generalized Calabi-Yau manifold with generalized Calabi-Yau structure φ of constant type k , and let T be a l -dimensional compact torus which acts on M in a Hamiltonian way. That is, this is a Lie group action on M which preserves the generalized Calabi-Yau structure φ , and there is a smooth function $\mu : M \longrightarrow \mathfrak{t}^*$ which satisfies

- (1) μ is T -equivariant, and
- (2) $\xi_M - \sqrt{-1}d\mu^\xi$ lies in E_φ for all $\xi \in \mathfrak{t}$, where ξ_M denotes the induced vector field on M and μ^ξ is the smooth function defined by $\mu^\xi(p) = \mu(p)(\xi)$.

We call the function μ the generalized moment map for the Hamiltonian action. Let dm be a natural volume form on M defined by

$$dm = \frac{(\sqrt{-1})^n}{2^{n-k}} \langle \varphi, \bar{\varphi} \rangle.$$

We proved in [6] that the pushforward measure $\mu_*(dm)$ on $\mathfrak{t}^*$ is absolutely continuous with respect to the Lebesgue measure $d\zeta$ on $\mathfrak{t}^*$, and gave the explicit formula of the density function.

Theorem 1.1 ([6], Theorem B.). *Let (M, φ) be a $2n$ -dimensional connected generalized Calabi-Yau manifold of constant type k , and suppose that compact l -torus T acts on M effectively and in a Hamiltonian way. In addition, We assume that the generalized moment map μ is proper. Then the pushforward $\mu_*(dm)$ of the natural volume form dm under μ is absolutely continuous with respect to the Lebesgue measure on $\mathfrak{t}^*$ and the Radon-Nikodym derivative f can be written by*

$$f(a) = \int_{M_a} dm_a = \text{vol}(M_a)$$

for each regular value $a \in \mathfrak{t}^*$ of μ , where dm_a denotes the measure defined by the natural volume form on the reduced space $M_a = \mu^{-1}(a)/T$.

The main purpose of this note is to prove the density function f is a piecewise polynomial.

Theorem 1.2. *Under the assumptions above, the function f is a polynomial (of degree $\leq n - k - l$) on each connected component of the set of regular values of the generalized moment map μ .*

The proof of Theorem 1.2 will be given in Section 2. Next using the Atiyah-Bott-Berline-Vergne localization theorem, we shall lead the localization formula for the oscillatory integral

$$\int_M e^{\sqrt{-1}\mu^\xi} dm, \quad \xi \in \mathfrak{t}$$

in the case that M is compact. The detailed notations and definitions is in the Section 3.

Theorem 1.3. *In addition to the assumptions in Theorem 1.2, we assume that M is compact. Then the localization theorem holds, i.e., the oscillatory integral is given by*

$$\int_M e^{\sqrt{-1}\mu^\xi} dm = \sum_{X \subset M^T} \frac{1}{2^{\dim X - l}} \int_X \frac{e^{i_X^* \mu^\xi} i_X^* (\varphi \wedge \sigma(\bar{\varphi}))}{e(NX)(\xi)},$$

for $\xi \in \mathfrak{t}$ such that $\beta_j(\xi) \neq 0$, where X is a connected component of the fixed point set M^T , $i_X : X \rightarrow M$ is the inclusion, and $e(NX)$ is the equivariant Euler class of the normal bundle of X .

2. PROOF OF THEOREM 1.2

Let M_{free} be the set on which T acts freely. Then since M is connected and T acts on M effectively, M_{free} is open dense subset of M and $M \setminus M_{\text{free}}$ has measure 0 (see [5], Appendix B). So we may assume T acts M freely, and it is sufficient to show that f is a polynomial of degree at most $n - l - k$. Let $p : M \rightarrow M/T$ be the natural projection. Since the generalized moment map $\mu : M \rightarrow \mathfrak{t}^*$ is T -invariant, there exists a smooth map

$$\psi : M/G \rightarrow \mathfrak{t}^*$$

such that

$$\mu = \psi \circ p.$$

Then $\mu^{-1}(a) = p^{-1}(\psi^{-1}(a))$ and $M_a = \psi^{-1}(a)$, for each regular value $a \in \mathfrak{t}^*$. Let $j_a : M_a \rightarrow M/G$ be the inclusion, so that we have the commutative relation

$$\pi \circ i_a = j_a \circ p_a,$$

where $i_a : \mu^{-1}(a) \rightarrow M$ is the inclusion, and $p_a : \mu^{-1}(a) \rightarrow M_a$ is the natural projection. Now we choose a connection on M and let θ^i be the curvature forms. Then the equivariant differential form

$$e^{2\sqrt{-1}\mu} \varphi \wedge \sigma(\bar{\varphi})$$

is closed, and using the Cartan map $\Omega_T(M) \rightarrow \Omega(M/T)$,

$$e^{2\sqrt{-1}\psi_i \theta^i} \wedge \nu \wedge \sigma(\bar{\nu})$$

is closed form, where ν is the unique form on M/T with the property that

$$p^* \nu = \varphi_{\text{hor}},$$

the horizontal part of φ (the definition of the Cartan map is in [5], for instance). From the relation $\pi \circ i_a = j_a \circ \pi_a$, we obtain

$$j_a^* \nu = \varphi_a.$$

Let c denote the cohomology class of $e^{2\sqrt{-1}\psi_i\theta^i} \nu \wedge \sigma(\bar{\nu})$, then

$$\begin{aligned} [\varphi_a \wedge \sigma(\bar{\varphi}_a)] &= j_a^* [\nu \wedge \sigma(\bar{\nu})] \\ &= j_a^* [e^{-2\sqrt{-1}\psi_i\theta^i} \wedge (e^{2\sqrt{-1}\psi_i\theta^i} \wedge \nu \wedge \sigma(\bar{\nu}))] \\ &= j_a^* (e^{-2\sqrt{-1}a_i[\theta^i]} \wedge c). \end{aligned}$$

Since M_a is compact and oriented, the embedding $j_a : M_a \rightarrow M/T$ defines a homology class $[M_a] \in H_{2(n-l)}(M/T, \mathbb{Z})$. This depends smoothly on a , and being an integer class is thus independent of a . So let us fix an a_0 in the image of μ . Then we have $[M_{a_0}] = [M_a]$. Thus the integral can be calculated by

$$\begin{aligned} f(a) &= \langle [M_a], [dm_a] \rangle \\ &= \left\langle [M_{a_0}], \left[\frac{(\sqrt{-1})^{(n-l)}}{2^{n-l-k}} (\varphi_a \wedge \sigma(\bar{\varphi}_a))_{2(n-l)} \right] \right\rangle \\ &= \left\langle [M_{a_0}], \frac{(\sqrt{-1})^{(n-l)}}{2^{n-l-k}} (e^{-2\sqrt{-1}a_i[\theta^i]} \wedge c)_{2(n-l)} \right\rangle, \end{aligned}$$

where $\langle, \rangle$ is the pairing of a homology class and a cohomology class. This shows that f is a polynomial of degree at most $n - l - k$.

3. PROOF OF THEOREM 1.3

Suppose that M is compact. Then the number of subgroups of T which can occur as stabilizer groups of points of M is finite, and so there exists a finite number of weights $\beta_1, \dots, \beta_m$ with the property that if $\xi \in \mathfrak{t}$ satisfies

$$\beta_i(\xi) \neq 0, \quad i = 1, \dots, m$$

the corresponding vector field ξ_M on M is non-zero except at points of M^T (see [5] for example).

Consider a complex form $\varphi' \in \Omega^{ev/od} M \otimes \mathbb{C}$ defined by

$$\varphi' = \varphi_k + \frac{1}{2}\varphi_{k+2} + \frac{1}{4}\varphi_{k+4} + \dots,$$

where φ_s is the s -th degree component of φ . Then a complex equivariant differential form $e^{\sqrt{-1}\mu}\varphi' \wedge \sigma(\bar{\varphi}')$ is equivariant closed, and $(\varphi' \wedge \sigma(\bar{\varphi}'))_{2n} = dm$. So by applying the Atiyah-Bott-Berline-Vergne localization theorem (see for instance [1] and [5]), we have

$$\begin{aligned} \int_M e^{\sqrt{-1}\mu^\xi} dm &= \int_M (e^{\sqrt{-1}\mu}\varphi' \wedge \sigma(\bar{\varphi}'))(\xi) \\ &= \sum_{X \subset M^T} \int_X \frac{i_X^* (e^{\sqrt{-1}\mu}\varphi' \wedge \sigma(\bar{\varphi}'))(\xi)}{e(NX)(\xi)} \\ &= \sum_{X \subset M^T} \frac{1}{2^{(\dim X - l)}} \int_X \frac{e^{i_X^* \mu^\xi} i_X^* (\varphi \wedge \sigma(\bar{\varphi}))}{e(NX)(\xi)}, \end{aligned}$$

for $\xi \in \mathfrak{t}$ satisfying $\beta_i(\xi) \neq 0$ for all $i = 1, \dots, m$, where X is a connected component of the fixed point set M^T , $i_X : X \longrightarrow M$ is the inclusion, and $e(NX)$ is the equivariant Euler class of the normal bundle of X . This shows Theorem 1.3.

REFERENCES

- [1] M. F. Atiyah, and R. Bott, The moment map and equivariant cohomology, *Topology* 23 no. 1, 1-28, (1984)
- [2] J. Duistermaat and G. Heckman, On the variation in the cohomology of the symplectic form of the reduced phase space, *Invent. Math* 69, 259-268 (1982)
- [3] J. Duistermaat and G. Heckman, Addendum to "On the variation in the cohomology of the symplectic form of the reduced phase space.", *Invent. Math* 72, 153-158 (1983)
- [4] V. Guillemin, V. Ginzburg and Y. Karshon, Moment maps, cobordisms, and Hamiltonian group actions, *Mathematical surveys and monographs*, volume 98
- [5] V. W. Guillemin and S. Sternberg, *Supersymmetry and Equivariant de Rham Theory*, Springer, 1999
- [6] Y. Nitta, Reduction of generalized Calabi-Yau structures , [math.DG/0611341](https://arxiv.org/abs/math.DG/0611341)

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY. 1-1 MACHIKANEYAMA, TOYONAKA, OSAKA 560-0043 JAPAN.